

ATTACHMENT 1

	FISH CONTAMINANT LEVEL								
	Generic Example 1ppm			Portland Harbor Bass (whole body) 0.25 to 4.5 ppm (by river mile)			Portland Harbor Carp (whole body) 5.9 ppm (site wide (95%UCL mean))		
	Superfund process		Compared to bkgrd in breast milk ⁴	Superfund process		Compared to bkgrd in breast milk ⁴	Superfund process		Compared to bkgrd in breast milk ⁴
	Cancer Risk	HQ		Cancer Risk	HQ		Cancer Risk	HQ	
Breast-feeding Child (mother consumes 142 g/day)	2 X 10 ⁻³	3200 ⁽¹⁾	75 times bkgrd	5 X 10 ⁻⁴ to 9 X 10 ⁻³		19 to 338 times bkgrd	1 x 10 ⁻²		443 times bkgrd
		600 ⁽²⁾							
		2100⁽³⁾			525 to 9,450⁽³⁾			12,000⁽³⁾	
Adult Fish Consumer (consumption of 142 g/day)				4 X 10 ⁻⁴ to 8 X 10 ⁻³	30 to 500		1 x 10 ⁻²	656	
Child Fish Consumer (consumption of 60 g/day)				2 X 10 ⁻⁴ to 3 X 10 ⁻³	200 to 800		4 x 10 ⁻³	1000	

¹ Assume 1 year of breastfeeding and use EPA RfD

² Assume 1 year of breastfeeding, 6 years of resident fish consumption and use EPA RfD

³ Assume 1 year of breastfeeding and use ATSDR sub-chronic (2 weeks to 1 year) MRL. Recommended approach (in bold).

⁴ The background concentration of PCBs in breast milk assumed for the Housatonic River site was 0.32 mg/kg-lipid compared to the concentration of 24 mg/kg-lipid estimated for the generic example of 1 ppm PCBs in fish.